

Validations using the Cryogenic Frostpoint Hygrometer (CFH)

Holger Vömel

Cooperative Institute for Research in Environmental Sciences
University of Colorado

Acknowledgements

Henry Selkirk, NASA Ames Research Center

Jéssica Valverde Canossa, Universidad Nacional

Juan Valdés, Universidad Nacional

Students of the Universidad Nacional and Universidad de Costa Rica

Werner Stoltz, Instituto Meteorológico Nacional

Grace Peng, Aerospace Corporation

J. Andrés Diaz, Centro Nacional de Alta Tecnología

Fumio Hasebe, Hokkaido University

Masatomo Fujiwara, Hokkaido University

Masato Shiotani, Kyoto University


Cornelius Schiller, FZ Jülich

Overview


- Update on MLS water vapor validation with version 1.5
- Quicklook at MLS water vapor test version 2.1
- Comparison of Hygrometers
- Vaisala RS92 update

Cryogenic Frostpoint Hygrometer (CFH)

- Absolute measurement
- Vertical Range: surface to ~28 km
(surface to ~25 km on ascent)
- Uncertainty: troposphere: > 4% MR
stratosphere: ~ 9 %
- Microprocessor control
- Phase sensitive detector:
electronic sunlight filter
- Weight: ~ 400 gr
- Payloads carry ECC ozone
sonde and Vaisala RS80
- ~170 soundings so far




Satellite comparison




Validation of MLS version 1.5 water vapor


MLS Comparison: Arctic winter



MLS Comparison: Mid latitudes




Boulder, CO




Lauder, NZ

MLS Comparison: Tropics

Costa Rica : July





July 2005



July 2006



MLS Comparison: Tropics

Biak Indonesia: Jan 2006





MLS Comparison: Tropics

Costa Rica AVE Jan/Feb 2006





Comparison of versions 1.5 and 2.1


Sondakylä & Boulder example



Comparison with FIRS-2



Average difference from CFH: version 1.5 & 2.1




Mid and high latitudes
only


Comparison between CFH and aircraft

January/February 2006

WB57 and CFH profiles




Harvard Ly-alpha




JPL TDL

CFH Correlations





Harvard Ly-alpha




JPL TDL

CFH Correlation MayComm balloon borne TDL




CR-AVE and SOWER/Biak comparison



Average
CFH profile at
Costa Rica
 $10.0^{\circ}\text{N}, 84.1^{\circ}\text{W}$
4 Jan – 4 Mar 2006

Average
CFH profile at
Biak Indonesia
 $1.2^{\circ}\text{S}, 136.1^{\circ}\text{E}$
8-16 Jan 2006

Geophysica and CFH profiles




**CFH profiles at
Biak Indonesia
1.2°S, 136.1°E
8-16 Jan 2006**

**FISH total water
profiles at
Darwin, Australia
12°S, 131°E
4 Nov – 14 Dec 2005**


Data courtesy C. Schiller

Daytime comparisons between CFH and Vaisala RS92

Relative RH difference RS92 - CFH



Costa Rica Jul 2005



Beltsville, MD Jul 2006

Summary

- Stratospheric MLS water vapor:
Agreement within measurement uncertainty for both version 1.5 and 2.1
- Except for tropical tape recorder during boreal winter
- Tropospheric MLS water vapor:
Version 2.1 improves general shape in UT,
but still very dry and highly variable
- WB57 instruments are too wet compared to CFH;
no serious disagreements between CFH and other balloon or Geophysica instruments
- Vaisala RS92 relative humidity during daytime still up to 50% too dry (same as last year)

**Validation of AURA/MLS water vapor
observations using the Cryogenic
Frostpoint Hygrometer (CFH) and
comparisons with aircraft water vapor
instruments and Vaisala RS92**

Holger Vömel

Cooperative Institute for Research in Environmental Sciences
University of Colorado